

**Interferometric Astrometry of
the Detached White Dwarf - M Dwarf Binary Feige 24
Using *Hubble Space Telescope* Fine Guidance Sensor 3:
White Dwarf Radius and Component Mass Estimates ¹**

G. Fritz Benedict¹, Barbara E. McArthur¹, Otto G. Franz², L. H. Wasserman², E. Nelan⁵, J. Lee⁷,
L.W. Fredrick¹², W. H. Jefferys⁶, W. van Altena⁷, E. L. Robinson⁶, W. J. Spiesman¹,
P. J. Shelus¹, P.D. Hemenway⁸, R. L. Duncombe⁹, D. Story¹⁰, A. L. Whipple¹¹, and A. Bradley¹¹

Received _____; accepted _____

¹Based on observations made with the NASA/ESA Hubble Space Telescope, obtained at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555

¹McDonald Observatory, University of Texas, Austin, TX 78712

²Lowell Observatory, 1400 West Mars hill Rd., Flagstaff, AZ 86001

⁵Space Telescope Science Institute, 3700 San Martin Dr., Baltimore, MD 21218

⁶Astronomy Dept., University of Texas, Austin, TX 78712

⁷Astronomy Dept., Yale University, PO Box 208101, New Haven, CT 06520

⁸Oceanography, University of Rhode Island, Kingston, RI 02881

⁹Aerospace Engineering, University of Texas, Austin, TX 78712

¹⁰ Jackson and Tull, Aerospace Engineering Division 7375 Executive Place, Suite 200, Seabrook, Md. 20706

¹¹Allied-Signal Aerospace, PO Box 91, Annapolis Junction, MD 20701

¹²Astronomy Dept., University of Virginia, PO Box 3818, Charlottesville, VA 22903

ABSTRACT

With *HST* FGS 3 we have determined a parallax for the white dwarf - M dwarf interacting binary, Feige 24. The white dwarf (DA) component has an effective temperature, $T_{eff} \sim 56,000K$. A weighted average with past parallax determinations ($\pi_{abs} = 14.6 \pm 0.4$ milliseconds of arc) narrows the range of possible radius values, compared to past estimates. We obtain $R_{DA} = 0.0185 \pm 0.0008 R_{\odot}$ with uncertainty in the temperature and bolometric correction the dominant contributors to the error. FGS photometry provides a light curve entirely consistent with reflection effects. A recently refined model Mass-Luminosity Relation (Baraffe et al. 1998) for low mass stars provides a mass estimate for the M dwarf companion, $\mathcal{M}_{dM} = 0.37 \pm 0.20 \mathcal{M}_{\odot}$, where the mass range is due to metallicity and age uncertainties. Radial velocities from Vennes and Thorstensen (1994) provide a mass ratio from which we obtain $\mathcal{M}_{DA} = 0.49^{+0.19}_{-0.05} \mathcal{M}_{\odot}$. Independently, our radius and recent $\log g$ determinations yield $0.44 < \mathcal{M}_{DA} < 0.47 \mathcal{M}_{\odot}$. In each case the minimum DA mass is that derived by Vennes & Thorstensen from their radial velocities and Keplerian circular orbits with $i \leq 90$ deg. Locating Feige 24 on an $\mathcal{M} - R$ plane suggests a carbon core.

Our radius and these mass estimates yield a γ_{grav} inconsistent with that derived by Vennes & Thorstensen. We speculate on the nature of a third component whose existence would resolve the discrepancy.

Subject headings: astrometry — stars: individual (Feige 24) — stars: distances — stars: binary — stars: white dwarfs — stars: late-type

1. Introduction

Feige 24 (= PG 0232+035 = HIP 12031) is DA white dwarf, red dwarf (M1-2V) (Liebert & Margon 1977) binary ($P = 4.23$ days, Vennes & Thorstensen 1994=VT94) that is described as the prototypical post-common envelope detached system with a low probability of becoming a Cataclysmic Variable (CV) within a Hubble time (King et al. 1994 and Marks, 1994). This object was selected for our *HST* parallax program because a directly measured distance could reduce the uncertainty of the radius of one of the hottest white dwarfs. Since the instigation of this program and the selection of targets over 15 years ago, at least two other groups have measured a parallax for Feige 24; USNO-Flagstaff (Dahn et al., 1988), and *HIPPARCOS* (Perryman et al., 1997 and Vaucclair et al., 1997). We outlined the results of a preliminary analysis in Benedict et al., 2000. Here we discuss our analysis and final results in detail.

Provencal et al. 1998 presented radii derived from *HIPPARCOS* parallaxes for 21 white dwarfs. In most cases, the dominating error term for the white dwarf radii was the parallax uncertainty. Our parallax of Feige 24, while slow in coming, has provided a fractional parallax uncertainty, $\frac{\Delta\pi}{\pi}$, similar to those in the Provencal et al. 1998 study, but for a much hotter, more distant object.

We time-tag our data with a modified Julian Date, $MJD = JD - 2400000.5$. We abbreviate millisecond of arc, mas; white dwarf, DA; and M dwarf, dM, throughout.

2. The Astrometry

Our astrometric observations were obtained with Fine Guidance Sensor 3 (FGS 3), a two-axis, white-light interferometer aboard *HST*. Bradley et al., 1991 provide an overview of the FGS 3 instrument and Benedict et al., (1999) describe the astrometric capabilities of FGS 3 and typical data acquisition and reduction strategies.

We use the term ‘pickle’ to describe the field of regard of the FGS. The instantaneous field of view of FGS 3 is a 5×5 arcsec square aperture. Figure 1 shows a finding chart for Feige 24 and our astrometric reference stars in the FGS 3 pickle as observed on 08 Aug 1997. Note the less than ideal placement of the primary science target with respect to the reference frame. The placement of Feige 24, at one side of the distribution of reference stars, seems to have produced no adverse astrometric or photometric effects.

2.1. The Astrometric Reference Frame

Table 3 provides a list of the observation epochs. Our data reduction and calibration procedures are described in Benedict et al., (1999) and McArthur et al., 1999. We obtained a total of 71 successful measurements of our reference stars during eight ‘observing runs’. For each of these eight observation sets we determine the scale and rotation relative to the sky, using a GaussFit (Jefferys et al., 1988) model. The orientation of the observation sets is obtained from ground-based astrometry (USNO-A2.0, Monet 1998) with uncertainties in the field orientation $\pm 0^\circ 12$.

Having only 8 observation sets and four reference stars precludes us from our usual practice (Benedict et al., (1999)) of constraining the proper motions and parallaxes to sum to zero ($\Sigma\mu = 0$ and $\Sigma\pi = 0$) for the entire reference frame. From a series of solutions we determined that only reference star ref-3 has a statistically significant proper motion and parallax. So, we constrain $\mu = 0$ and $\pi = 0$ for reference stars ref-2, -4 and -5.

We conclude from histograms (Figure 2) of the reference star residuals that we have obtained a per-observation precision of ~ 1 mas. The resulting reference frame ‘catalog’ (Table 2) was determined with final errors $< \sigma_\xi > = 0.5$ and $< \sigma_\eta > = 0.6$ mas.

To determine if there might be unmodeled, but eventually correctable, systematic effects at the 1 mas level, we plotted the Feige 24 reference frame X and Y residuals against a number of spacecraft, instrumental, and astronomical parameters. These included X, Y position within the pickle; radial distance from the pickle center; reference star V magnitude and B-V color; and epoch of observation. We saw no trends, other than the expected increase in positional uncertainty with reference star magnitude.

2.2. Modeling the Parallax and Proper Motion of Feige 24

Spectroscopy of the reference frame stars obtained from the WIYN ¹ and an estimate of color excess, E(B-V), from Burstein & Heiles 1982. (Table 2) shows that the colors of the reference stars and our science target differ, with $\Delta(B-V) \sim -1$. Therefore, we apply the differential correction for lateral color discussed in Benedict et al., (1999) to the Feige 24 observations and obtain a parallax relative to our reference frame,

¹The WIYN Observatory is a joint facility of the University of Wisconsin-Madison, Indiana University, Yale University, and the National Optical Astronomy Observatories.

$\pi_{rel} = 13.8 \pm 0.4$ mas. The proper motion relative to the four astrometric reference stars is listed in Table 4.

Franz et al (1998) and Benedict et al., (1999) have demonstrated 1 mas astrometric precision for FGS 3. Table 3 presents our Feige 24 astrometric residuals obtained from the parallax and proper motion model. Histograms of these residuals are characterized by $\sigma_x = 1.0$ and $\sigma_y = 1.2$ mas. This was slightly larger than expected. To investigate whether or not the larger residuals could be attributed to Feige 24, Figure 3 presents the residuals phased to the VT94 orbital period, $P = 4.23160^d$, with $T_0 = HJD\ 2448578.3973$. We find no significant trends in the astrometric residuals. In particular, there is no correlation with the two distinct *HST* orientations required by the pointing constraints discussed in Benedict et al., (1999). With any reasonable masses for the DA and dM components, a binary system at this distance, having this period, could exhibit maximum reflex motion at the 0.5 mas level. This null detection does not place very useful upper limits on the component masses.

Because our parallax for Feige 24 is determined with respect to the reference frame stars which have their own parallaxes, we must apply a correction from relative to absolute parallax. The WIYN spectroscopy and the estimated color excess (See Table 2) indicate a reference frame with an average parallax of $\langle \pi \rangle_{ref} = 0.9 \pm 0.4$ mas. where the error is based on the dispersion of the individual spectrophotometric parallaxes. To check our correction to absolute, we compare it to those used in the Yale Parallax Catalog (YPC95, van Altena, Hoffleit, & Lee, Section 3.2). From YPC95, Fig. 2, the Feige 24 galactic latitude, $b = -50^\circ 3$ and average magnitude for the reference frame, $\langle V_{ref} \rangle = 13.4$, we obtain a correction to absolute of 1.9 mas. Rather than use a galactic model-dependent correction, we adopt the spectroscopically derived $\langle \pi \rangle_{ref} = 0.9 \pm 0.4$ mas. Applying this correction results in an absolute parallax of $\pi_{abs} = +14.7 \pm 0.6$ mas, where the error has equal contributions from the HST FGS observations and the correction to absolute parallax. Finally, we note that our proper motion is smaller than either the HIPPARCOS or USNO values, where the HIPPARCOS value is an absolute proper motion while the USNO and the HST values are relative to their respective reference frame proper motions. If our reference stars are a representative statistical sample of the parent population, then based on the data in Table III in van Altena (1974), we expect a statistical uncertainty in the mean value of the correction to absolute proper motion (not applied here) of ± 6 mas y^{-1} .

We compare our absolute parallax to previous work in Table 4 and in Figure 4. We adopt for the remainder of this paper the weighted average absolute parallax, $\langle \pi_{abs} \rangle = 14.6 \pm 0.4$ mas, shown as a horizontal dashed line in Figure 4. Weights used are $1/\sigma^2$.

Lutz & Kelker (1973) show that for a uniform distribution of stars, the measured trigonometric parallaxes are strongly biased towards the observer (i.e., too large), rendering inferred distances and luminosities too small. This bias is proportional to $(\sigma_\pi/\pi)^2$. Using a space density determined for the CV RW Tri (McArthur et al. 1999), and presuming that Feige 24 is a member of that same class of object (binaries containing white dwarfs), we determine an LK correction of -0.01 ± 0.01 magnitudes. Correcting our distance modulus, we obtain $m - M = 4.17 \pm 0.11$.

2.3. Kinematic Age of the Feige 24 System

From the VT94 systemic radial velocity and either our proper motions or those from HIPPARCOS (Table 4) we derive the space velocity of Feige 24, 67 ± 1 km s⁻¹. The velocity component perpendicular to the galactic plane, W , is -37 km s⁻¹. Our new parallax places the star 53 parsecs below the Sun or 61 parsecs below the galactic plane. An object this far below the galactic plane and continuing to move further away from the plane so swiftly is more characteristic of a ‘thick disk’ than a thin disk object (c.f. Thejll et al., 1997). Feige 24, if truly a Pop I object, has a space velocity 3.5 times the young disk velocity dispersion. These data suggest that Feige 24 formed prior to the formation of the galactic disk, although subsequent evolution of the DA component is likely quite recent. This may be an instance of past mass transfer in an intermediate Population II object.

3. Astrophysics of the Feige 24 System

We discuss the consequences of a more precisely determined parallax, calculating some astrophysically relevant parameters for the DA and dM components. These are collected in Table 5. Our goals are the radius and mass of the DA component. We first calculate a radius, then estimate the time since the DA formation event. Component masses have been estimated by VT94. We will revisit this issue later. That we do not substantially improve the mass uncertainty motivates a future direct measurement of the component separation. This one measurement would yield precise masses. A series of measurements would provide individual orbits, possibly illuminating past and future component interactions.

3.1. Estimating the DA Radius

To estimate the DA radius we require an intrinsic luminosity. From Landolt 1983 we obtain a system total magnitude, $V_{tot} = 12.41 \pm 0.01$. The magnitude of the white dwarf is critical and difficult to obtain, because the M dwarf always contributes flux. Holberg et al., 1986 derive $V_{DA} = 12.56 \pm 0.05$ using IUE spectra. They ratio Feige 24 with other hot DA; G191 B2B, GD246 and HZ43. From the DA magnitude and total magnitude we obtain $V_{dM} = 14.63 \pm 0.05$ and $\Delta V = 2.07$. We assume an $A_V = 0$ for Feige 24 at $d = 69$ pc, consistent with our adopted $A_V = 0.09$ for the reference frame at an average distance $d = 1600$ pc (Table 2). The LK bias-corrected distance modulus ($m - M = 4.17 \pm 0.11$) then yields absolute magnitudes $M_V = 10.46 \pm 0.12$ for the red dwarf companion and $M_V = 8.39 \pm 0.12$ for the DA.

A recently determined temperature of the Feige 24 DA, taking into account non-LTE and heavy element effects (Barstow Hubeny & Holberg 1998), is $T_{eff}^{DA} = 56,370 \pm 1,000K$. This temperature yields a radius via differential comparison with the sun. This procedure requires a bolometric magnitude, hence, a bolometric correction. We could adopt the bolometric correction, B.C. = -4.88, generated by Bergeron et al. (1995) from a pure Hydrogen, $\log g = 8$ DA model convolved with a V bandpass. But, Feige 24 is neither $\log g = 8$ nor pure H.

Flower (1996) provides bolometric corrections for normal stars up to $T_{eff} \sim 54,000K$. From Flower (1996), figure 4, the relationship between $\log T_{eff}$ and B.C. is linear for $T_{eff} > 25,000K$. Hotter stars lie on the Rayleigh-Jeans tail of the blackbody curve, where flux is roughly proportional to T_{eff} , not T_{eff}^4 . A small linear extrapolation yields B.C. = -4.82 $\pm$ 0.06 for the Feige 24 DA. The B.C. error comes from the uncertainty in T_{eff}^{DA} . Because a DA with some heavy elements in its atmosphere radiates more like a hot normal star than a pure H DA, we choose the Flower correction rather than the model correction. We are also encouraged by the near equality of the B.C. from observation and theory.

We obtain a DA bolometric luminosity $M_{bol}^{DA} = M_V + B.C. = 3.57 \pm 0.13$. R_{DA} follows from the expression

$$M_{bol}^{\odot} - M_{bol}^{DA} = 10 \log(T_{eff}^{DA} / T_{eff}^{\odot}) + 5 \log(R_{DA} / R_{\odot}) \quad (1)$$

where we assume for the Sun $M_{bol}^{\odot} = +4.75$ and $T_{eff}^{\odot} = 5800K$. We find $R_{DA} = 0.0180 \pm 0.0013R_{\odot}$, following the error analysis of Provencal et al. 1998. The primary sources of error for this radius are the bolometric correction and the T_{eff}^{DA} .

A second approach to deriving R_{DA} involves the V-band average flux, H_V , discussed in Bergeron et al.

(1995). They list H_V^{DA} as a function of temperature for, again, the pure Hydrogen, $\log g = 8$ model. If we can determine an $H_V^\odot$, we can derive R_{DA} from

$$R_{DA} = (H_V^\odot / H_V^{DA}) 10^{-0.4(M_V^{DA} - M_V^\odot)} \quad (2)$$

where $M_V^{DA} = 8.39 \pm 0.12$ comes from our parallax and $M_V^\odot = 4.82$ is assumed. We obtain $H_V^\odot$ by convolving the Bessel (1990) V band response with the solar spectral distribution listed in Allen (1973). We calculate $H_V^\odot = 6.771 \times 10^5 \text{ ergs cm}^{-2} \text{ s}^{-1} \text{ \AA}^{-1} \text{ str}^{-1}$. We obtain for $T_{eff} = 56,370 K$ an $R_{DA} = 0.0188 \pm 0.0010$.

A weighted average of the two independent determinations provides $R_{DA} = 0.0185 \pm 0.0008 R_\odot$, where the error is certainly underestimated due to unknown systematic effects. Parallax is no longer a significant source of error for the radius determination. Comparing with the results presented in Provencal et al. 1998, figure 7, we find Feige 24 to have a radius larger than any other white dwarf.

With a temperature $T_{eff} \sim 56,000$, the time since the DA formation event is unlikely to be longer than 1.5 My. This conclusion is drawn from the DA cooling tracks as function of mass calculated by M. Wood, detailed in Sion 1999, fig. 7. These models also indicate that the DA mass must satisfy $\mathcal{M}_{DA} \geq 0.4 \mathcal{M}_\odot$ to remain near this lofty T_{eff} for longer than 3×10^5 y.

3.2. Estimating The White Dwarf Mass

Before estimating $\mathcal{M}_{DA}$ we review the VT94 minimum component masses from their radial velocities and the Kepler relation for total system mass, separation, and period. Then, we estimate the DA mass using two different approaches. We first attempt to determine the most likely dM mass. The VT94 radial velocity amplitude ratio then provides the DA mass. The second, independent mass estimate follows from our derived radius along with the DA atmospheric parameter, $\log g$, obtained through spectroscopy. Our DA mass estimate will differ little from VT94, and, if better, is so only by virtue of more recent dM models and DA atmospheric parameters.

3.2.1. Minimum Component Masses from Binary Radial Velocities

The system total lower mass limit can be set by the VT94 radial velocities and the Kepler relation for mass, separation, and period. VT 94 give us the velocities along each component orbit, the fact that each orbit is circular (from the pure sine wave fits to the velocity curves), and the period, the time

it takes to travel around each orbit. Assuming an edge-on system ($i = 90^\circ$), one that can produce the full vector amount of radial velocity amplitude measured by VT94, the minimum system mass is $\mathcal{M}_{tot} = 0.73\mathcal{M}_\odot$. From the VT94 mass ratio, $\mathcal{M}_{dM}/\mathcal{M}_{DA} = 0.63 \pm 0.04$, we obtain the DA mass limit, $\mathcal{M}_{DA \text{ Kepler}} \geq 0.44\mathcal{M}_\odot$, and the dM mass limit, $\mathcal{M}_{dM \text{ Kepler}} \geq 0.26\mathcal{M}_\odot$. No smaller masses can produce the observed radial velocities for orbits of these known sizes. At $d = 68.5$ pc an edge-on system with minimum mass would separate the components by 672 microarcsec, or $9.9R_\odot$.

3.2.2. Inclination from the Light Curve

VT 94 find $H\alpha$ equivalent width variations that phase with the orbital period. These show a maximum at $\phi = 0.5$. Photometric variations of Feige 24 might be detectable, because the photometric capabilities of FGS 3 approach a precision of 0.002 magnitude (Benedict et al., 1998). Figure 5 shows the flat-fielded counts and the corresponding differential instrumental magnitudes as well as a sin wave fit with amplitude and phase as free parameters. There is a clear photometric signature with a peak-to-peak amplitude 0.028 magnitude, showing maximum system brightness at phase $\phi = 0.58 \pm 0.09$. Given the sparse coverage, this phase at maximum is not surprisingly different from the $H\alpha$ equivalent width maximum seen at $\phi = 0.5$.

A likely mechanism for producing the single-peaked orbital light curve is heating of the dM star by the white dwarf (the reflection effect). As the dM star orbits the white dwarf, its heated face is alternately more or less visible, increasing and decreasing the observed flux from Feige 24 once per orbit. To test this hypothesis we calculated model light curves using an updated version of the light curve synthesis program described by Zhang et al. (1986). We initially adopted $T_{eff} = 56,370$ K and $R = 0.0185 R_\odot$ for the white dwarf, $T_{eff} = 3800$ K and $R = 0.52 R_\odot$ for the M1-2V star, and 4.8×10^{-2} AU ($10.3R_\odot$) for the separation of their centers of mass, and then adjusted the temperature of the dM star so that it contributed 13.5% of the V flux from the system. The peak-to-peak amplitudes of the resulting model light curves are a function of orbital inclination, topping out at ~ 0.025 mag for $i = 90^\circ$, and can easily be made to agree in amplitude and shape with the observed light curve.

This photometric behavior is entirely consistent with reflection effects (c.f. Robinson et al. 2000). We find that the quality of the observed light curve is, however, inadequate to improve the parameters of the system, particularly the inclination. We have not sufficiently sampled the expected flat section of the light curve (near $\phi = 0$). Nevertheless, these results do provide quantitative evidence that (1) the orbital light curve is caused by heating and (2) the heating is consistent with the radius and temperature we have

derived for the white dwarf – a useful external check on our results.

3.2.3. DA Mass from the M Dwarf

The dM absolute magnitude ($M_V = 10.46 \pm 0.12$) implies a spectral type M2V (Henry, Kirkpatrick, & Simons 1994), consistent with Liebert & Margon 1977. The absolute magnitude of an M dwarf star depends not only on mass, but also on age (evolutionary stage) and chemical composition. Baraffe et al., 1998 have produced a grid of models, varying metallicity, $[M/H]$ and helium abundance, Y . We plot in Figure 6 their Mass-Luminosity curves for dwarfs with ages 10My and 10Gy, with $[M/H] = 0$ and 10Gy with $[M/H] = -0.5$, all with solar helium abundance. The complete grid of Baraffe et al. models shows that M dwarfs in the mass range $0.175 \leq \mathcal{M}_{dM} \leq 0.43\mathcal{M}_\odot$ with $-0.5 < [M/H] < 0$ have $M_V = 10.46$ at some time in their evolution from 10My to 10Gy.

The dM mass now depends on metallicity and how quickly an M dwarf of a given mass decreases in brightness. Figure 7 shows the dependence of brightness on mass, age, and metallicity. These Baraffe et al. models indicate that solar metallicity stars with higher mass remain near $M_V = 10.46$ far longer than low mass stars. However, kinematically, Feige 24 is more likely to be old and of lower than solar metallicity than young and of normal metallicity. First adopting the 10Gy model, $[M/H] = 0$, and calculated absolute magnitude, we estimate the dM star mass $\mathcal{M}_{dM} = 0.43 \pm 0.08\mathcal{M}_\odot$, because that mass remains at $M_V = 10.46$ for a larger fraction of the total lifetime than any other. However, if we accept the kinematical suggestion of allegiance to a thick disk population, then $[M/H] < 0$ is more likely. Assuming $[M/H] = -0.5$ results in a dM star mass $\mathcal{M}_{dM} = 0.185 \pm 0.08\mathcal{M}_\odot$.

Radial velocities from VT94 (dM from Kitt Peak, DA from IUE) provide the velocity amplitude ratio, $K_{DA}/K_{dM} = 0.63 \pm 0.04 = \mathcal{M}_{dM}/\mathcal{M}_{DA}$. From the total possible dM mass range, $0.185 < \mathcal{M}_{dM} < 0.43\mathcal{M}_\odot$, and the mass ratio we derive a DA mass range, $0.29 < \mathcal{M}_{DA} < 0.68\mathcal{M}_\odot$. Applying the limit, $\mathcal{M}_{DA \text{ Kepler}} \geq 0.44\mathcal{M}_\odot$, we obtain $0.44 < \mathcal{M}_{DA} < 0.68\mathcal{M}_\odot$. Keplerian lower limits argue for a dM star mass $0.26 < \mathcal{M}_{dM} < 0.43\mathcal{M}_\odot$, a range consistent with a metallicity slightly less than solar and an age in excess of 0.3 Gy (Figure 7).

3.2.4. DA Mass from Atmospheric Parameters

The dM star does not provide a particularly precise DA mass estimate. If one knows the surface gravity, g , and the radius, R , the mass can be obtained through

$$M = gR^2/G \quad (3)$$

where G is the gravitational constant. The quantity $\log g$ comes from analysis of the line profiles in spectra. Recent determinations include: Marsh et al. 1997, $\log g = 7.53 \pm 0.09$; Kidder 1991, $\log g = 7.45 \pm 0.51$; Vennes et al., 1997, $\log g = 7.2 \pm 0.07$; Finley Koester & Basri, 1997, $\log g = 7.17 \pm 0.15$; and Barstow Hubeny & Holberg 1998, $\log g = 7.36 \pm 0.12$. The full range of the measures and Equation 3 yield the range of mass values $0.21 \leq \mathcal{M}_{DA} \leq 0.47\mathcal{M}_{\odot}$. Applying the limit, $\mathcal{M}_{DA \text{ Kepler}} \geq 0.44\mathcal{M}_{\odot}$ eliminates nearly all of these mass determinations. In this case our radius and the Kepler limit indicate that $\log g$ should be at the high end of these measures.

3.3. The White Dwarf Composition

We next place Feige 24 on a white dwarf mass-radius diagram (Figure 8). We plot our two independently determined mass ranges against our adopted radius, $R_{DA} = 0.0185 \pm 0.0008R_{\odot}$. We represent the radius error by the two horizontal long-short dashed lines. The top thick horizontal bar shows the $\mathcal{M}_{DA}$ determined from atmospheric parameters. Only the largest $\log g$ at the largest radius produces masses in excess of the Keplerian limit. The thick bar at $R_{DA} = 0.0185R_{\odot}$ indicates the $\mathcal{M}_{DA}$ range derived through the dM mass estimates. For this determination the mass error bars indicate the range of ages and $[M/H]$ discussed in section 3.2.2. For any dM older than 1-2 Gy the lower masses are associated with lower metallicity. The vertical bold dotted line shows the lowest possible $\mathcal{M}_{DA}$ that can produce the observed VT 94 radial velocity amplitudes for an edge-on orientation of this binary system. We also plot several values of $\log g$ (dashed) and γ_{grav} (thin solid). The wide grey curves in Figure 8 are C and He DA models from Vennes et al., 1995. While uncertain, a carbon core DA seems more likely than a pure He core DA.

4. Discussion

While our estimated dM and DA masses differ little from VT94, our DA radius differs substantially. VT94 note the difference between their minimum radius, $R_{DA} = 0.028R_{\odot}$, and that predicted by the Dahn

et al., 1988 parallax. This discrepancy is exacerbated by the two new parallax determinations (*HST* and *HIPPARCOS*), folded into our weighted average parallax.

VT94 derive a DA gravitational redshift, $\gamma_{grav} = 8.7 \pm 2 \text{ km s}^{-1}$ from the measured mean velocities for the dM and DA. Combined with our $R_{DA} = 0.0185 R_{\odot}$, this γ_{grav} suggests a forbidden DA mass, $\mathcal{M}_{DA} \sim 0.3 \mathcal{M}_{\odot}$. Reducing the mass of the DA component could reconcile the VT94 $\log g$ and γ_{grav} with our radius.

We speculate that a third component in the Feige 24 system, a low-mass companion to the DA star, could preserve the total system mass and lower the DA mass. If all components are coplanar, the VT94 DA radial velocities apply strict limits to this reconciliation, because too high a mass for component C would show up as large residuals. We estimate from the scatter that a radial velocity amplitude of $\pm 10 \text{ km s}^{-1}$ could ‘hide’ in the VT94 DA radial velocity measurements. Stellar dynamics applies yet another constraint. Holman & Wiegert (1999) parameterize the stability of tertiary companions as a function of stellar component A and B mass function, $\mu = \mathcal{M}_A / (\mathcal{M}_A + \mathcal{M}_B)$, and AB binary orbit ellipticity, e . With $e=0$ and $\mu = 0.39$ we find (from their table 3) that component C must have an orbital semi-major axis less than 0.3 times that of AB.

Insisting that $\mathcal{M}_{DA} = 0.30 \mathcal{M}_{\odot}$ (this mass - with our radius - would produce the upper limit VT94 $\gamma_{grav} = 10.7 \text{ km s}^{-1}$) requires $\mathcal{M}_C = 0.14 \mathcal{M}_{\odot}$ ($\mathcal{M}_A + \mathcal{M}_C = 0.44 \mathcal{M}_{\odot}$). To hide the C component from the radial velocity technique requires a very low AC inclination, nearly face-on. However, non-coplanarity reduces the size of the stable AC semi-major axis even further (Weigert & Holman, 1997; Pendleton & Black 1983). As an example suppose component C must have an orbital semi-major axis of 0.1 or less that of AB to insure stability. An AC period, $P = 0.18^d$ (4.3^h), and $i = 6^\circ$ would produce a radial velocity signature of about $\pm 10 \text{ km s}^{-1}$. Finally, the Mass-Luminosity Relation of Henry et al. (1999) would predict $M_V^C = 14.0$, hence, $V_C \sim 17.2$, likely undetectable in any of the spectra analyzed for radial velocities. Have we built a new CV, one that should evidence mass transfer and all the associated phenomena? A recent review of CVs (Beuermann, 1999) indicates that the putative component C ($\mathcal{M}_C = 0.14 \mathcal{M}_{\odot}$) would have to orbit much closer ($P \sim 1.5^h$) to the DA primary before filling its Roche lobe and producing the characteristic signature of a CV.

Finally we note that our radius differs little from that derived by VT94 from the only trigonometric parallax then available (Dahn et al., 1988). The unresolved inconsistency between radii (derived from direct parallaxes) and surface gravities (derived from minimum mass and those radii) illuminates the need for

high angular resolution observations and direct mass determinations.

The Feige 24 DA mass will rest on an age- and metallicity-dependent lower main sequence Mass-Luminosity Relationship or still uncertain $\log g$ measurements until the component separations are measured directly. Resolving the inconsistencies between the DA mass estimates (involving dM stellar models and uncertain temperatures, $\log g$, and bolometric corrections) requires astrometry, both to further reduce the parallax uncertainty, and, more importantly, to spatially resolve this system. Astrometrically derived orbital parameters will provide unambiguous and precise mass determinations for both components. They may also offer insight regarding past and future component interactions.

This system and dozens more like it are ideal targets for the Space Interferometry Mission (<http://sim.jpl.nasa.gov>). Feige 24, at a distance of 69 pc with $P = 4.23^d$, has a total component separation on order 700 microarcsec. The component orbits are much larger than the expected SIM measurement limits. Because shortward of 700 nm 70-80% of the system flux is contributed by the DA (Thorstensen et al. 1978), the wide SIM bandpass and spectral resolution should allow measurement of positions, magnitudes, and colors for both components, even with $\Delta V \sim 2$.

Once launched SIM will provide crucial astrometry for this and similar systems at ten times the distance (determined by target magnitude, not astrometric precision). SIM measurements of this system along with many other binaries will provide data with which to create an age- and metallicity-dependent Mass-Luminosity Relationship of exquisite accuracy.

5. Conclusions

1. The weighted average of three independent parallax measurements yields a distance to the dM + DA binary Feige 24 with $\sigma_D/D = 2.8\%$. $D = 68.4^{+2.0}_{-1.9}$ pc.
2. We estimate the radius of the DA component using two methods. The first requires either a model-dependent bolometric correction, or one that derives from hot, normal stars. The second utilizes a model-dependent V-band average flux, H_V . The two results agree within their errors and yield a weighted average $R_{DA} = 0.0185 \pm 0.0008 R_\odot$, where the most significant contributions to the error are the uncertain T_{eff}^{DA} and B.C.. This radius is larger than any of the WD discussed in Provencal et al. 1998.
3. FGS photometry provides quantitative evidence that the orbital light curve is caused by heating of the dM component by the DA. That signature is consistent with the assumed temperature and the radius

we have derived for the white dwarf

4. The VT94 measured radial velocity amplitudes, amplitude ratios, and the assumption of Keplerian circular motion exclude $\mathcal{M}_{DA\ Kepler} < 0.44\mathcal{M}_{\odot}$ and $\mathcal{M}_{dM\ Kepler} < 0.26\mathcal{M}_{\odot}$.

5. We estimate the dM component mass, $0.26 < \mathcal{M}_{dM} < 0.43\mathcal{M}_{\odot}$, from the Baraffe et al., 1998 stellar evolution models, a lower limit from Keplerian circular orbits, and the VT 94 radial velocities. The upper range is due to unknown age and metallicity, [M/H]. A DA mass range ($0.44 < \mathcal{M}_{DA} < 0.68\mathcal{M}_{\odot}$) follows directly from the VT94 radial velocity amplitudes.

6. We determine $\mathcal{M}_{DA}$ from our R_{DA} and a rather wide range of spectroscopically determined $\log g$ values. This approach yields $0.44 \leq \mathcal{M}_{DA} \leq 0.47\mathcal{M}_{\odot}$, where again the lower limit is imposed by $\mathcal{M}_{DA\ Kepler} > 0.44\mathcal{M}_{\odot}$.

7. We plot these DA component mass ranges on the $\mathcal{M} - R$ plane. With the assistance of the hard lower mass limit and C and He DA $\mathcal{M} - R$ models from Vennes et al., 1995, we identify Feige 24 to have a carbon core. A pure He core DA seems less likely.

8. Noting that our radius and the minimum possible $\mathcal{M}_{DA}$ are inconsistent with the VT94 γ_{grav} , we explore the possibility of a tertiary component. A component C, orbiting a common center of mass with the DA, having a period in the range $1.5 < P < 5^h$ with the orbit plane nearly face-on, could reduce the DA mass to $\mathcal{M}_{DA} = 0.30\mathcal{M}_{\odot}$ and not produce any observational evidence.

8. SIM will be able to measure the orbits of each known component and provide directly measured dynamical masses for both. Orbit size and precise shape may provide information on the nature of past and future interactions between the two components. SIM would also detect a tertiary, if present.

This research has made use of NASA’s Astrophysics Data System Abstract Service and the SIMBAD Stellar Database inquiry and retrieval system. We gratefully acknowledged web access to the astrometry and photometry in USNO-A2.0, provided by the United States Naval Observatory, Flagstaff Station. Support for this work was provided by NASA through grant GTO NAG5-1603 from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. We thank Don Winget and Sandi Catalán for discussions and draft paper reviews. Denise Taylor provided crucial scheduling assistance at the Space Telescope Science Institute. Travis Metcalfe kindly provided code for JD to HJD corrections. We thank an anonymous referee for suggestions that enhanced the clarity of our presentation, and for giving us the courage to speculate.

REFERENCES

- Allen, C. W. 1973, London: University of London, Athlone Press, 3rd ed.,
- Baraffe, I., Chabrier, G., Allard, F. & Hauschildt, P. H. 1998, *A&A*, 337, 403
- Barstow, M. A., Hubeny, I. & Holberg, J. B. 1998, *MNRAS*, 299, 520
- Beuermann, K., 1999. To appear in *New Astronomy Reviews* (astro-ph/9909141)
- Bradley, A., Abramowicz-Reed, L., Story, D., Benedict, G. & Jefferys, W. 1991, *PASP*, 103, 317
- Benedict, G. Fritz, McArthur, Barbara, Nelan, E., Story, D., Whipple, A. L., Shelus, P. J., Jefferys, W. H., Hemenway, P. D., Franz, Otto G., Wasserman, L. H., Duncombe, R. L., Van Altena, W., Fredrick, L. W., 1998, *AJ*, 116, 429
- Benedict, G. F., McArthur, B., Chappell, D. W., Nelan, E., Jefferys, W. H., Van Altena, W., Lee, J., Cornell, D., Shelus, P. J., Hemenway, P. D., Franz, O. G., Wasserman, L. H., Duncombe, R. L., Story, D., Whipple, A., & Fredrick, L. W. 1999, *AJ*, 118, 1086
- Benedict, G. F., McArthur, B., Franz, O. G., Wasserman, L. H., Nelan, E., Fredrick, L. W., Van Altena, W., Lee, J., Jefferys, W. H., Shelus, P. J., Hemenway, P. D., Duncombe, R. L., Story, D., & Whipple, A., 2000, to appear in *ASP Proceedings "Working on the Fringe"*
- Bergeron, P., Wesemael, F. & Beauchamp, A. 1995, *PASP*, 107, 1047
- Bessell, M. S. 1990, *PASP*, 102, 1181
- Burstein, D. & Heiles, C. 1982, *AJ*, 87, 1165
- Dahn, C. C., Harrington, R. S., Kallarakal, V. V., Guetter, H. H., Luginbuhl, C. B., Riepe, B. Y., Walker, R. L., Pier, J. R., Vrba, F. J., Monet, D. G. & Ables, H. D. 1988, *AJ*, 95, 237
- Finley, David S., Koester, Detlev & Basri, Gibor 1997, *ApJ*, 488, 375
- Flower, P. J. 1996, *ApJ*, 469, 355

- Franz, O. G., Henry, T. J., Wasserman, L. H., Benedict, G. F., Ianna, P. A., Kirkpatrick, J. D., McCarthy, D. W., Jr., Bradley, A. J., Duncombe, R. L., Fredrick, L. W., Hemenway, P. D., Jefferys, W. H., McArthur, B. E., Nelan, E. P., Shelus, P. J., Story, D. B., Van Altena, W. F. & Whipple, A. L. 1998, AJ, 116, 1432
- Henry, T. J., Franz, O. G., Wasserman, L. H., Benedict, G. F., Shelus, P.J., Ianna, P.A., Kirkpatrick, J. D., & McCarthy, D. W. 1999, ApJ, 512, 864
- Henry, T. J., Kirkpatrick, J. D. & Simons, D. A. 1994, AJ, 108, 1437
- Holberg, J. B., Basile, J. & Wesemael, F. 1986, ApJ, 306, 629
- Holman, M. & Wiegert, P., AJ, 117, 621
- Jefferys, W., Fitzpatrick, J., and McArthur, B. 1988, Celest. Mech. 41, 39.
- Kidder, Kenneth Mark 1991, Ph.D. Thesis, University of Arizona
- King, A.R., Kolb, U., De Kool, M. & Ritter, H. 1994, MNRAS, 269, 907
- Landolt, A. U. 1983, AJ, 88, 439
- Liebert, J. & Margon, B. 1977, ApJ, 216, 18
- Lutz, T. E. & Kelker, D. H. 1973, PASP, 85, 573
- Marsh, M. C., et al. 1997, MNRAS, 286, 369
- McArthur, B. E., Benedict, G. F., Lee, J., Lu, C. -L., Van Altena, W. F., Deliyannis, C. P., Girard, T., Fredrick, L. W., Nelan, E., Duncombe, R. L., Hemenway, P. D., Jefferys, W. H., Shelus, P. J., Franz, O. G. & Wasserman, L. H. 1999, ApJ, 520, L59
- Marks, P. B., 1994, in Cataclysmic Variables, eds. Bianchini, A., Della Valle, M. and Orio, M., Kluwer, p. 535
- Monet, D. G. 1998, American Astronomical Society Meeting, 193, 112.003
- Nelan, E. "FGS Instrument Handbook ver 8", 1999, STScI
- Pendleton, Y. J. & Black, D. C. 1983, AJ, 88, 1415

- Perryman, M. A. C., Lindegren, L., Kovalevsky, J., Høg, E., Bastian, U., Bernacca, P. L., Crézé, M., Donati, F., Grenon, M., Van Leeuwen, F., Van Der Marel, H., Mignard, F., Murray, C. A., Le Poole, R. S., Schrijver, H., Turon, C., Arenou, F., Froeschlé, M. & Petersen, C. S. 1997, *A&A*, 323, L49
- Provencal, J. L., Shipman, H. L., Høg, E. & Thejll, P. 1998, *ApJ*, 494, 759
- Robinson, E. L., Welsh, W. F. & Young, P. 2000, to appear in *X-ray Astronomy 1999: Stellar Endpoints, AGN and the X-ray Background* Conference Proceedings, Bologna, Italy
- Sion, E. M. 1999, *PASP*, 111, 532
- Thejll, P., Flynn, C., Williamson, R. & Saffer, R. 1997, *A&A*, 317, 689
- Thorstensen, J. R., Charles, P. A., Bowyer, S. & Margon, B. 1978, *ApJ*, 223, 260
- Van Altena, W.F., Lee, J.T. & Hoffleit, E.D. 1995, New Haven, CT: Yale University Observatory, 4th ed., completely revised and enlarged
- van Altena, W., 1974, *AJ*, 79, 826
- Vauclair, G., Schmidt, H., Koester, D. & Allard, N. 1997, *A&A*, 325, 1055
- Vennes, S. & Thorstensen, J. R. 1994, *AJ*, 108, 1881 (VT94)
- Vennes, S., Fontaine, G. & Brassard, P. 1995, *A&A*, 296, 117
- Vennes, S. , Thejll, P. A., Galvan, R. G. & Dupuis, J. 1997, *ApJ*, 480, 714
- Wiegert, P. A. & Holman, M. J. 1997, *AJ*, 113, 1445
- Wood, M.A., 1990, Ph.D. Thesis. University of Texas at Austin
- Zhang, E., Robinson, E. L., & Nather, R. E. 1986, *ApJ*, 305, 740

Table 1. Feige 24 Reference Frame: Astrometry

ID	V	ξ (arcsec)	η (arcsec)	μ_X (arcsec y ⁻¹)	μ_Y (arcsec y ⁻¹)	π (arcsec)
ref-2*	11.59	0.0±0.0004	0.0±0.0004	0	0	0
ref-3	13.38	-118.9943±0.0004	50.4968±0.0004	0.0157±0.0004	- 0.0014±0.00050	-0.0006±0.0003
ref-4	14.82	27.3168±0.0010	63.5709±0.0010	- 0	0	0
ref-5	13.66	-144.4703±0.0004	86.9851±0.0005	0	0	0

*RA, Dec = 288.087967, 2.898281 (J2000)

Table 2. Feige 24 and its Reference Frame: Stellar Parameters

ID	V ^a	$B-V$ ^b	SpT	M_V	E(B-V) ^c	A_V	m-M	D(pc)	π_{abs} (mas)
ref-2	11.59	1.00	G9 III	0.75	0.03	0.093	10.84	960	0.6
ref-3	13.38	0.69	G3 V	4.8	0.03	0.093	8.58	410	1.9
ref-4	14.82	0.61	G0 V	4.4	0.03	0.093	10.42	935	0.8
ref-5	13.66	0.63	F9 III	1.2	0.03	0.093	12.46	2540	0.3
Feige 24	12.41 ^d	-0.20 ^d							

^afrom FGS PMT measures calibrated as per Nelan et al.1999

^bfrom $B-V = f(\text{Sp.T.}) + E(B-V)$

^cfrom Burstein & Heiles 1982

^dfrom Landolt 1983

Table 3. *HST* Observations of Feige 24 and Astrometric Residuals

Obs. Set	MJD	X residual	Y residual
1	49930.92188	0.0002	0.0008
1	49930.9375	0.0000	-0.0012
1	49930.94531	-0.0003	-0.0004
2	49936.88672	0.0005	-0.0002
2	49936.90234	-0.0002	-0.0003
2	49936.91016	0.0006	0.0001
3	50102.09375	-0.0014	0.0008
3	50102.10938	0.0001	-0.0002
3	50102.11719	0.0004	-0.0013
4	50109.06641	-0.0008	0.0007
4	50109.07813	-0.0001	0.0002
4	50109.08594	0.0007	0.0001
5	50669.78125	-0.0001	0.0010
5	50669.79688	0.0009	0.0016
5	50669.80469	-0.0010	-0.0004
6	50678.92188	-0.0003	-0.0004
6	50678.9375	0.0003	0.0006
6	50678.94531	-0.0016	0.0010
7	50819.28125	0.0014	-0.0003
7	50819.28906	0.0005	-0.0008
7	50819.29688	0.0003	0.0001
8	50821.29688	0.0006	-0.0009
8	50821.30469	-0.0005	-0.0008
8	50821.3125	-0.0007	0.0006

Table 4. Feige 24 Parallax, Proper Motion, and Radial Velocity

Parameter	Value	
<i>HST</i> study duration	2.4 y	
number of observation sets	8	
Number of ref. stars	4	
ref. stars $\langle V \rangle$	13.4	
ref. stars $\langle B - V \rangle$	0.7	
<i>HST</i> Relative Parallax	13.8 ± 0.4 mas	
corr to absolute	0.9 ± 0.7 mas	
<i>HST</i> Absolute Parallax	14.7 ± 0.6 mas	
<i>HIPPARCOS</i> Absolute Parallax	13.4 ± 3.6 mas	
USNO Absolute Parallax	13.5 ± 2.9 mas	
<i>HST</i> Proper Motion (μ)	71.1 ± 0.6 mas y ⁻¹	
in p.a.	83°6	
<i>HIPPARCOS</i> μ	85.8 ± 5 mas y ⁻¹	
in p.a.	84°2	
USNO μ	78.4 ± 1.9 mas y ⁻¹	
in p.a.	88°4	
Weighted Average Absolute Parallax	14.6 ± 0.4 mas	
m-M (LK bias corrected)	4.17 ± 0.11	
System Radial Velocity, γ	$+ 62.0 \pm 1.4$ km s ⁻¹	VT94
Galactocentric z velocity, W	-37 ± 1.5 km s ⁻¹	γ & <i>HST</i> or <i>HIPPARCOS</i> μ

Table 5. Feige 24 Astrophysical Quantities

Parameter	Value	Source
V_{tot}	12.41 ± 0.01	Landolt 1983
$B - V$	-0.20 ± 0.01	Landolt 1983
V_{DA}	12.56 ± 0.05	Holberg et al., 1986
V_{dM}	14.63 ± 0.05	V_{tot} & V_{DA}
A_V	0	reference frame $< B - V >$ & Sp.T. (Table 2)
m-M (LK bias corrected)	4.17 ± 0.11	this paper
dM M_V	10.46 ± 0.12	m-M
dM Sp. T.	M2V	dM M_V & Henry, Kirkpatrick, & Simons 1994
$\mathcal{M}_{dM}$	$0.29 - 0.43 \mathcal{M}_{\odot}$	dM M_V , Baraffe et al., 1998, $\mathcal{M}_{DA}$ <i>Kepler</i>
DA M_V	8.39 ± 0.12	m-M
DA B.C.	-4.82 ± 0.06	Flower, 1996
M_{bol}^{DA}	3.57 ± 0.13	$= DA M_V + B.C.$
T_{eff}^{DA}	$56,370 \pm 1,000 K$	Barstow Hubeny & Holberg 1998
R_{DA}	$0.0185 \pm 0.0008 R_{\odot}$	this paper
$\mathcal{M}_{dM}/\mathcal{M}_{DA}$	0.63 ± 0.04	K_{DA}/K_{dM} VT94
$\mathcal{M}_{DA}$	$0.49^{+0.19}_{-0.05} \mathcal{M}_{\odot}$	$\mathcal{M}_{dM}$, K_{DA}/K_{dM} & $\mathcal{M}_{DA}$ <i>Kepler</i>
$\mathcal{M}_{DA}$	$0.44 - 0.47 \mathcal{M}_{\odot}$	$\log g$ & $\mathcal{M}_{DA}$ <i>Kepler</i>




Fig. 1.— Location of reference stars within the FGS 3 field of regard on 08 Aug 1997. Note the less than ideal placement of the primary science target with respect to the reference frame.




Fig. 2.— Histograms of x and y residuals obtained from modeling the Feige 24 reference frame to obtain scale, orientation, and offset parameters. Distributions are fit with gaussians.




Fig. 3.— Astrometric residuals in RA (X) and Dec (Y) phased to the VT94 orbit ($P = 4.23160^d$ and $T_0 = JD2448578.3973$). Boxes and open circles denote the two *HST* orientations, which seem to have no effect on the astrometric residuals. Dashed lines are best-fit sine waves constrained to the VT94 period.




Fig. 4.— Absolute parallax determinations for Feige 24. We compare *HST*, *HIPPARCOS*, and USNO (Dahn et al., 1988). Error bars are 1- σ . The horizontal dashed line gives the weighted average absolute parallax, $\langle \pi_{\text{abs}} \rangle$.




Fig. 5.— Flat fielded intensity (the filter (F583W) has a bandpass centered on 583 nm, with 234 nm FWHM) and differential instrumental magnitudes phased to the VT94 orbit ($P = 4.23160^d$ and $T_0 = JD2448578.3973$). Boxes and open circles denote the two *HST* orientations, which seem to have no effect on the photometry. The dashed line is a best-fit sine wave constrained to the VT94 period.




Fig. 6.— M dwarf absolute magnitude as a function of mass from the stellar evolution models of Baraffe et al., 1998. A wide range of masses, ages, and metallicities can result in the derived dM absolute magnitude, $M_V = 10.46$ (horizontal dashed line). Note that at a given mass, a low metallicity ($[M/H] = -0.5$) star is always brighter than a high metallicity ($[M/H] = 0.0$) star.




Fig. 7.— The time variation of absolute magnitude for M dwarfs of various masses taken from the stellar evolution models of Baraffe et al., 1998. Empty symbols are for solar metallicity; filled for low metallicity ($[M/H]=-0.5$). Note that the higher mass stars remain near the computed M dwarf absolute magnitude, $M_V = 10.46$ (dashed line), far longer than the low mass stars. A low metallicity star with $\mathcal{M}_{dM} = 0.175\mathcal{M}_{\odot}$ is brighter than a solar metallicity star with $\mathcal{M}_{dM} = 0.40\mathcal{M}_{\odot}$.




Fig. 8.— Feige 24 mass and radius on a DA mass-radius map. The vertical bold dotted line shows the lowest possible $\mathcal{M}_{DA}$ from keplerian considerations. The radius error is represented by the top and bottom long-short dashed lines. The top thick horizontal bar shows the $\mathcal{M}_{DA}$ determined from atmospheric parameters. Only at the largest radius (lowest temperature) and largest $\log g$ do we obtain a DA mass in excess of the keplerian limit. The bottom thick horizontal bar at $R_{DA} = 0.0185R_{\odot}$ indicates the $\mathcal{M}_{DA}$ range derived through the dM. We also plot several values of $\log g$ (dashed) and γ_{grav} (thin solid). The grey wide solid lines are C and He DA models from Vennes et al., 1995. A C core DA is somewhat more likely than an He core DA.